

Influence of ventilation use and occupant behaviour on surface microorganisms in contemporary social housing

Sharpe T.¹, McGill G,^{1*} Dancer SJ,^{2,3} King M-F,⁴ Fletcher L,⁴ Noakes CJ.⁴

Affiliations:

¹ Department of Architecture, University of Strathclyde, Scotland, UK

² Dept. of Microbiology, Hairmyres Hospital, NHS Lanarkshire, Scotland, UK

³ School of Applied Sciences, Edinburgh Napier University, Scotland, UK

⁴ Water, Public Health and Environmental Engineering Group, School of Civil Engineering, University of Leeds, England, UK

*Correspondence to: grainne.mcgill@strath.ac.uk

Running title: Effect of ventilation on home surface microbes

Keywords: Ventilation; Community; Microbiology; Bacteria; Fungi; Surfaces

Supplementary Information

Table 3: Cultivable Gram-negative bacteria identified from eight sampled sites in Scottish homes.

Most isolates were *Pantoea* spp. or *Pantoea agglomerans*

Also found:

Acinetobacter baumannii (also *lwoffii*; *radioresistens*; *ursingii*; *haemolyticus*)

Sphingomonas paucimobilis; *Pseudomonas putida* (also *fluorescens*)

Lerclercia adecarboxylata

Paracoccus yeei

Klebsiella pneumoniae

Enterobacter cloacae

Roseomonas gilardii

Aeromonas sobria

Serratia liquefaciens

Brevundimonas spp.

Cronobacter sakazakii

Moraxella spp.